#### **BIBLIOGRAPHIC INFORMATION SYSTEM**

Journal Full Title: Journal of Biomedical Research & Environmental Sciences Journal NLM Abbreviation: J Biomed Res Environ Sci Journal Website Link: https://www.jelsciences.com Journal ISSN: 2766-2276 Category: Multidisciplinary Subject Areas: Medicine Group, Biology Group, General, Environmental Sciences **Topics Summation: 128 Issue Regularity: Monthly** Review Process: Double Blind Time to Publication: 21 Days Indexing catalog: Visit here Publication fee catalog: Visit here

#### • **DOI:** 10.37871 (CrossRef)

Plagiarism detection software: iThenticate

Managing entity: USA

Language: English

Research work collecting capability: Worldwide

Organized by: SciRes Literature LLC

**License:** Open Access by Journal of Biomedical Research & Environmental Sciences is licensed under a Creative Commons Attribution 4.0 International License. Based on a work at SciRes Literature LLC.

**IndexCopernicus** 

ICV 2020:

53.77

Manuscript should be submitted in Word Document (.doc or .docx) through

### **Online Submission**

form or can be mailed to support@jelsciences.com

Tision: Journal of Biomedical Research & Environmental Sciences main aim is to enhance the importance of science and technology to the scientific community and also to provide an equal opportunity to seek and share ideas to all our researchers and scientists without any barriers to develop their career and helping in their development of discovering the world.



# Minimize Risks of COVID-19 Infection

### Sunil J Wimalawansa\*

Professor of Medicine, Endocrinology, & Nutrition; Cardiometabolic and Endocrine Institute, NJ, USA

### **EDITORIAL**

The first wave of COVID-19 spread across the globe, rapidly during the first half of the year 2020 [1]. Since August 2020, the second wave of COVID-19 has been rampaging across most countries. A third wave may likely occur during the late spring of 2021. These in part coincide with the annual winter flu season in countries located in northern and later, the southern most latitudes, during their respective winter periods. One exception is the summertime peak in the middle-east when most people, near totally avoid exposure to sunlight due to extremely hot weather. During these vulnerable periods, body stores and the serum 25(OH)vitamin D [25(OH)D)] concentration reduce; in parallel, the immune system weakens, thus increasing the risks for respiratory viral illnesses, including COVID-19.

Many tropical countries reporting fewer infections, which is mostly due to conducting significantly less (i.e., less than 5%) PCR testing per million population compared to the west). In addition, inevitable exposure to sunlight of people living in equatorial countries prevent the occurrence of severe vitamin D deficiency. The combination is responsible for low number of PCR positivity and COVID-related deaths reported in tropical countries, not due to lockdowns, curfew, or healthcare.

## Why certain groups of people develop complications and dying from COVID-19?

While the reported prevalence and the deaths from COVID-19 are lowest in most tropical counties, it is not necessary the high temperature and high humidity alone responsible for this [2]. People in these regions get exposed to a reasonable amount of Ultraviolet B Rays (UVB) from the sunlight, facilitating the generation of vitamin D through the skin [3]. Consequently, they can maintain a reasonably active immune system preventing COVID-19 infections and complications [4]. Because food contains insufficient amounts of vitamin D, one must depend on UVB exposure and/or taking adequate doses of supplements to maintain optimum serum 25(OH)D concentration to maintain good health [5].

Having low vitamin D concentration weakens the immunity thereby increasing the risk of contracting COVID-19 [6], and associated complications and deaths [7-9]. While the first wave of COVID-19 could have affected anyone, the elderly and those in nursing homes had the brunt of developing complications, the use of ICU facilities, and [10] and most of the deaths. Those who were most vulnerable (elderly with chronic comorbidities and very low serum 25(OH)D concentrations) had the worst outcomes. The second wave, however, was also affecting the middle age and an increasing number of younger people who are relatively protected during the first wave of COVID-19. While the exact reason is unknown, human behavior generating higher viral loads are likely to have had an impact [11]. The latter was in part due to over-relaxation of restrictions by governments and the younger people take it for granted and began excessive socialization without taking public

#### \*Corresponding author

Sunil J Wimalawansa, Professor of Medicine, 661 Darmody Avenue, North Brunswick, NJ 08902

Tel: +1(732)940 0811 E-mail: suniljw@hotmail.com

DOI: 10.37871/jbres1174

Submitted: 18 December 2020

Accepted: 19 December 2020

Published: 19 December 2020

Copyright: © 2020 Wimalawansa SJ. Distributed under Creative Commons CC-BY 4.0

#### OPEN ACCESS

Subject: Medicine Group

Topic & Subtopic(s): Nutritional Disorders; Nutrition

#### Keywords

- Disease prevention
- Coronavirus
- Hypovitaminosis D
- Infection
- Inflammation
- Immunity
- Micronutrients
- Pandemic
- SARS-CoV-2

VOLUME: 1 ISSUE: 8



health precautions [12]. While certain strains demonstrating increased virulency, most others either remain the same or, becoming less virulent, perhaps due to ongoing mutations of the viruses and the remaining non-infected people having a higher degree of immunity.

## Steps that individuals could take to prevent COVID-19

There are several straightforward steps that individuals and families should take to reduce the risk of contracting COVID-19. These can be categories into three groups: avoiding the viral infection and the viral load (protecting self), preventing infecting others and enhance / strengthen immunity (protecting others), enabling to fight against the virus (boost the immunity and the use of vaccines). The first group includes wearing effective facemask such as N95, each time one goes out of home or from the office, washing hands with soap and water when return home and back to the workplace, avoiding crowded situations, minimize the use of public transportation and travelling (Figure 1).

The second group includes avoiding infecting others. This includes, avoid getting exposed to high-risk persons (who have a fever, coughing, or sneezing), and keeping a minimum of a four-meter distance from elder parents and relatives from younger children and grandchildren, who could be asymptomatic carriers. Nevertheless, the relationships between the elders and the youngsters should be strengthened using social media, video catting, and virtual visits, several times a day. Similarly, staying at home when has signs and/or symptoms of respiratory tract illness, strictly wearing effective facemasks, use a clean tissue or bend elbow if one has to cough or sneeze, and seeking appropriate medical assistances appropriate (Telemedicine first, calling a local emergency healthcare number for advice before rushing to a hospital).

The third group of actin necessary is to strengthen the immune system using natural means. Despite the use of macro-nutritional remedies, syrups, and herbal mixtures that sell, and propagated by television, radio, the social medial, and newspaper advertisement virtually has little or no effect on strengthening the immune system, that is capable of overpowering COVID-19. These are no better than a placebo. Nevertheless, there are other rational steps one can take to boost the immune system. Vitamin D also protects against all acute, viral respiratory infections [13-15]. These include maintaining micronutrient adequacy, especially vitamin D, vitamin C, K2, zinc and selenium, essential fatty acids such as omega-3, and magnesium [16].

Additive beneficial effects of these facilitate maintaining a robust immune system. The body also needs adequate rest and sleep as well as reasonable daily physical activity to maintain not only the mental and physical well-being but also the optimal function of the immune system [17]. Figure 1 illustrates the common modes of actions one should adhere to during the current COVID-19 pandemic to reduce the risks of acquiring the infection.

## Steps that governments should take to prevent COVID-19

Governments and administrations must have a comprehensive common-sense plan for controlling the transmission of COVID-19 based on the biology and the known modes of spread of the virus. In addition to the standard public health measures such as mandating the use of facemasks, adhere to maintaining social distance as COVID-19 is predominantly an airborne disease, expand detection systems using community-based PCR, rapid



Figure 1 Illustrates the three most important items that one could adhere to minimize COVID-19 infection.

### 😂 Liferature

Subject Area(s): NUTRITIONAL DISORDERS NUTRITION

antigen test, and antibody testing, and effective contact tracing using trained public health personnel.

Those who are exposed and even found positive do not necessarily need hospitalization unless they are developing any one of the known complications of COVID. Others can be provided with safe isolation at home under supervision. However, those who do not have such accommodation must be provided humane quarantining at government, health department (not military) run centres, free of charge to minimize cross infections. Meanwhile, it is possible that COVID-virus might mutate to generate a super-bug and the third wave could become out of control and most vaccines might lose their efficacy from late 2021 against COVID. Besides, governments must have an active, realtime surveillance system based on Geographic Information Systems (GIS) to identify locations of persons positive for COVID-19. Such would allow early identification of outbreaks' and COVID clusters, and delineate disease location with higher concentration to protect the most vulnerable people. The latter include nursing homes, disability centres, and elderly homes, and early location of hot spots thus, enabling them to take proactive actions to protect people. The latter include provision of high doses of vitamin D to high-risk communities to rapidly stimulate their immune system, enabling subduing COVID-19.

#### The need for a humane approach

During the process of controlling COVID, one must strike a balance between the virus spread versus the provision of basic human needs to the populous, the safety of the elders and children, etc. Broader and sensible evaluation of the need for opening of schools and universities versus continuing online schooling, safe transportation of children to school and people to work sites, and implementation of community-based disease prevention measures. Since some of the community spreads are occurring through workplaces and schools, re-structuring work routines and shifts, logistics, encouraging and provision of facilities to work from home for non-essential staff, provide safe transportation and workplace, the safety of staff, availability of free COVID-19 screening facility at work to all employees at free of charge, etc., are crucial for safety, and early identifications of developing cluster and managing them.

# Importance of early identification and isolation to stop the spread of COVID-19

The mentioned measures would allow prompt early identifications of positive cases and trace contacts, enabling prompt control of community viral spread. Locking down and curfews are not the answers to controlling COVID-19 [18]. Besides, all ports of entry, such as airports and seaports must have rapid screening facilities (e.g., rapid antigen test and others) for COVID-19 together with affirmative and humane quarantining. In addition, the public must be educated to prevent those who are ill and having a

temperature from travelling out of home, especially using public transportation. None of these will work in isolation in the absence of public education (but not through fearmongering, misleading, and threats) together with empowering them to take care of themselves and local communities.

### **SUMMATION**

At present, the only proven medical or nutritional intervention that reduces COVID-19 infection and reduces its complications and deaths is sun-exposure and higher doses of vitamin D supplementation [19]. Those who are at high risk, following exposure to a person with COVID-19 or a person with symptoms of COVID-19 infection likely to benefit from an upfront loading oral dose, such as between 100,000 and 400,000 IU given as a stat dose or over a few days could mitigate the illness swiftly. The same approach should be considered in all COVID-19 clinical trials conducted using vitamin D as an intervention. Another rescue means following an exposure, is to take 50,000 IU daily for 4 to 7 days, followed with a maintenance dose of 4,000 IU/day. The same approach should be considered in all COVID-19 clinical trials conducted using vitamin D as an intervention. This is especially important as those people with chronic diseases, the elderly, and those in nursing homes, who are known to have a high prevalence of severe vitamin D deficiency and are at high risk.

While the mentioned are highly cost-effective approaches, no government is yet to approve vitamin D as a preventative, adjunct, or a therapeutic agent. The latter is in part due to negative propaganda by the big-pharma, vaccine manufacturers, and their surrogates spreading false and disparaging information on vitamin D and bolstering and exaggerating the putative benefits from expensive antiviral agents and vaccines. Collectively, this propaganda misleads decision-makers and thus, preventing them from taking affirmative right actions to benefit the public.

#### References

- Wimalawansa SJ. Global epidemic of coronavirus-COVID-19: What can we do to minimize risks? European J Biomed & Pharma Sci. 2020;7: 432-438.
- Auler AC, Cássaro FAM, da Silva VO, Pires LF. Evidence that high temperatures and intermediate relative humidity might favor the spread of COVID-19 in tropical climate: A case study for the most affected Brazilian cities. Sci Total Environ. 2020 Aug 10;729:139090. doi: 10.1016/j.scitotenv.2020.139090. Epub 2020 Apr 28. PMID: 32388137; PMCID: PMC7194794.
- Miraglia Del Giudice M, Indolfi C, Strisciuglio C. Vitamin D: Immunomodulatory Aspects. J Clin Gastroenterol. 2018 Nov/Dec;52 Suppl 1, Proceedings from the 9th Probiotics, Prebiotics and New Foods, Nutraceuticals and Botanicals for Nutrition & Human and Microbiota Health Meeting, held in Rome, Italy from September 10 to 12, 2017:S86-S88. doi: 10.1097/MCG.000000000001112. PMID: 30300262.
- Ianevski A, Zusinaite E, Shtaida N, Kallio-Kokko H, Valkonen M, Kantele A, Telling K, Lutsar I, Letjuka P, Metelitsa N, Oksenych V, Dumpis U, Vitkauskiene A, Stašaitis K, Öhrmalm C, Bondeson K, Bergqvist A, Cox RJ, Tenson T, Merits A, Kainov DE. Low Temperature and Low UV Indexes Correlated with Peaks of Influenza Virus Activity in Northern Europe during 2010:2018. Viruses. 2019 Mar 1;11(3):207. doi: 10.3390/ v11030207. PMID: 30832226; PMCID: PMC6466003.

## Eliferature

- 5. Wimalawansa SJ. Biology of Vitamin D. J steroids Horm Sci. 2019;198:1-8.
- Wimalawansa SJ. Achieving population vitamin D sufficiency will markedly reduce healthcare costs. EJBPS. 2020;7:136-141.
- Trovas G, Tournis S. Vitamin D and COVID-19. Hormones (Athens). 2020 Jul 14:1–2. doi: 10.1007/s42000-020-00231-9. Epub ahead of print. PMID: 32666357; PMCID: PMC7360122.
- Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. Aging Clin Exp Res. 2020 Jul;32(7):1195-1198. doi: 10.1007/s40520-020-01570-8. Epub 2020 May 6. PMID: 32377965; PMCID: PMC7202265.
- D'Avolio A, Avataneo V, Manca A, Cusato J, De Nicolò A, Lucchini R, Keller F, Cantù M. 25-Hydroxyvitamin D Concentrations Are Lower in Patients with Positive PCR for SARS-CoV-2. Nutrients. 2020 May 9;12(5):1359. doi: 10.3390/nu12051359. PMID: 32397511; PMCID: PMC7285131.
- Cao X. COVID-19: Immunopathology and its implications for therapy. Nat Rev Immunol. 2020 May;20(5):269-270. doi: 10.1038/s41577-020-0308-3. PMID: 32273594; PMCID: PMC7143200.
- Neher RA, Dyrdak R, Druelle V, Hodcroft EB, Albert J. Potential impact of seasonal forcing on a SARS-CoV-2 pandemic. Swiss Med Wkly. 2020 Mar 16;150:w20224. doi: 10.4414/smw.2020.20224. PMID: 32176808.
- 12. Wimalawansa SJ. COVID-19: Evolution and prevention. Trends Telemed E-Health. 2020;2:1-5. https://bit.ly/3p4106k
- Morris SK, Pell LG, Rahman MZ, Dimitris MC, Mahmud A, Islam MM, Ahmed T, Pullenayegum E, Kashem T, Shanta SS, Gubbay J, Papp E, Science M, Zlotkin S, Roth DE. Maternal vitamin D supplementation during pregnancy and lactation to prevent acute respiratory infections in infancy in Dhaka, Bangladesh (MDARI trial): protocol for a prospective cohort study nested within a randomized controlled trial. BMC Pregnancy Childbirth. 2016 Oct 13;16(1):309. doi: 10.1186/s12884-016-1103-9. PMID: 27737646; PMCID: PMC5064894.

- Carpagnano GE, Di Lecce V, Quaranta VN, Zito A, Buonamico E, Capozza E, Palumbo A, Di Gioia G, Valerio VN, Resta O. Vitamin D deficiency as a predictor of poor prognosis in patients with acute respiratory failure due to COVID-19. J Endocrinol Invest. 2020 Aug 9:1–7. doi: 10.1007/s40618-020-01370-x. Epub ahead of print. PMID: 32772324; PMCID: PMC7415009.
- Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, Dubnov-Raz G, Esposito S, Ganmaa D, Ginde AA, Goodall EC, Grant CC, Griffiths CJ, Janssens W, Laaksi I, Manaseki-Holland S, Mauger D, Murdoch DR, Neale R, Rees JR, Simpson S Jr, Stelmach I, Kumar GT, Urashima M, Camargo CA Jr. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. BMJ. 2017 Feb 15;356:i6583. doi: 10.1136/bmj.i6583. PMID: 28202713; PMCID: PMC5310969.
- Wimalawansa SJ. Fighting against COVID-19: Boosting the immunity with micronutrients, stress reduction, physical activity, and vitamin D. Nutrition and Food Science Journal (Sci Literature). 2020;3:1-4.
- Wimalawansa SJ, Reducing risks from COVID-19: Cost-effective ways of strengthening individual's and the population immunity with vitamin D. J Endocrinol Sci, 2020. 2(2): p. -13. https://bit.ly/38cMz97
- Killeen GF, Kiware SS. Why lockdown? Why national unity? Why global solidarity? Simplified arithmetic tools for decision-makers, health professionals, journalists and the general public to explore containment options for the 2019 novel coronavirus. Infect Dis Model. 2020 Jul 3;5:442-458. doi: 10.1016/j.idm.2020.06.006. PMID: 32691016; PMCID: PMC7342051.
- Merzon E, Tworowski D, Gorohovski A, Vinker S, Golan Cohen A, Green I, Frenkel-Morgenstern M. Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study. FEBS J. 2020 Sep;287(17):3693-3702. doi: 10.1111/febs.15495. Epub 2020 Aug 28. PMID: 32700398; PMCID: PMC7404739.