

BIBLIOGRAPHIC INFORMATION SYSTEM

Journal Full Title: Journal of Biomedical Research & Environmental Sciences

Journal NLM Abbreviation: J Biomed Res Environ Sci

Journal Website Link: <https://www.jelsciences.com>

Journal ISSN: 2766-2276

Category: Multidisciplinary

Subject Areas: Medicine Group, Biology Group, General, Environmental Sciences

Topics Summation: 128

Issue Regularity: Monthly

Review Process type: Double Blind

Time to Publication: 7-14 Days

Indexing catalog: [Visit here](#)

Publication fee catalog: [Visit here](#)

DOI: 10.37871 ([CrossRef](#))

Plagiarism detection software: [iThenticate](#)

Managing entity: USA

Language: English

Research work collecting capability: Worldwide

Organized by: [SciRes Literature LLC](#)

License: Open Access by Journal of Biomedical Research & Environmental Sciences is licensed under a Creative Commons Attribution 4.0 International License. Based on a work at SciRes Literature LLC.

Manuscript should be submitted in Word Document (.doc or .docx) through **Online Submission** form or can be mailed to support@jelsciences.com

Vision

Journal of Biomedical Research & Environmental Sciences main aim is to enhance the importance of science and technology to the scientific community and also to provide an equal opportunity to seek and share ideas to all our researchers and scientists without any barriers to develop their career and helping in their development of discovering the world.

REVIEW ARTICLE

Human Immunodeficiency Virus (HIV) Drug Resistance: A Global Narrative Review

Maureen Nkandu Phiri and Steward Mudenda*

Department of Pharmacy, School of Health Sciences, University of Zambia, P.O. Box 50110, Lusaka, Zambia

ABSTRACT

Background: Antiretroviral Therapy (ART) has significantly improved Human Immunodeficiency Virus (HIV) patients' survival rates. However, the emergence of HIV Drug Resistance (HIVDR) has markedly reduced the effectiveness of Antiretroviral Therapy (ART).

Aim: This narrative review was conducted to review published studies on HIV drug resistance and its consequences.

Materials and methods: A literature search for this narrative review was carried out and the following databases were used PubMed, Google Scholar, and The Lancet. The cited articles were published from 1999 to 2021. The keywords used in the search of literature included 'Antiretroviral therapy', 'resistance', and 'Human Immunodeficiency Virus drug resistance', 'HIV', 'HIV drug resistance', 'HIV vaccines', and the Boolean word 'AND'.

Results: There is a high prevalence of HIV drug resistance globally that has been associated with some factors such as older age, non-adherence to treatment, long treatment duration, lower cell count and high viral load. HIV drug resistance may lead to treatment failure, prolongation of the time required to achieve viral suppression and leads to increased mortality. Increasing access to viral load monitoring can help mitigate HIV drug resistance.

Conclusion: HIV drug resistance is a global threat to public health and has been associated with increased morbidity and mortality. Therefore, there is a need for more research to be carried out and various strategies like the use of antiretrovirals with a high genetic barrier to resistance need to be put in place to prevent further spread resistance. HIVDR must be monitored frequently taking into consideration the geographic variability. There is an urgent need for the development of anti-HIV vaccines that will help to prevent further transmission and spread of HIV.

*Corresponding author

Steward Mudenda, Department of Pharmacy, School of Health Sciences, University of Zambia, P.O Box 50110, Lusaka, Zambia

Tel: +260-977-549-974

E-mail: freshsteward@gmail.com

DOI: 10.37871/jbres1323

Submitted: 15 September 2021

Accepted: 29 September 2021

Published: 30 September 2021

Copyright: © 2021 Phiri MN, et al. Distributed under Creative Commons CC-BY 4.0

OPEN ACCESS

Keywords

- HIV
- AIDs
- Antiretroviral therapy
- HIVDR
- HIV drug resistance
- HIV vaccines

INTRODUCTION

Human Immunodeficiency Virus (HIV) is a global public health issue that has led to increased morbidity and mortality [1-3]. The rapid expansion of Antiretroviral Therapy (ART) has significantly improved the prognosis of HIV positive patients, by reducing the morbidity and mortality rates in both developed and resource-limited countries [4-6]. The effectiveness of ART has been markedly reduced by the development of HIV Drug Resistance (HIVDR), which is a major factor in the failure of ART [7,8].

HIVDR occurs because of one or more changes in the genetic structure of HIV that affects the ability of a specific drug or combination of drugs to block the replication of the virus [9]. HIVDR is also observed when the ART fails to suppress the viral load [10,11]. Besides, HIVDR has been reported due to the widespread usage of ART to treat HIV [12]. HIVDR has become a significant barrier to reaching the Joint United Nations Programme on HIV and AIDS (UNAIDS) Fast-Track goal of ending Acquired

MEDICINE GROUP

HIV/AIDS

VOLUME: 2 ISSUE: 9



Immune Deficiency Syndrome (AIDS) by 2030 [13]. The threat of HIVDR is high, especially in sub-Saharan Africa and other low and middle-income countries where weak health systems and poor access make managing HIV more challenging [14,15]. The rise in HIVDR rates has mostly been against Nucleoside Reverse Transcriptase Inhibitors (NRTIs) and Non-nucleoside Reverse Transcriptase Inhibitors (NNRTIs) [16]. This is of particular concern because these drug classes are part of the first or/and second-line ART regimens.

Factors associated with the development of HIV drug resistance include; poor adherence, suboptimal ART doses, initiation of therapy late in the course of HIV infection, use of substandard antiretroviral regimens, lack of plasma viral load monitoring, the transmission of drug-resistant HIV, treatment interruptions, high replication rates of the virus, selective pressure caused by the Antiretrovirals (ARVs), and initial infection by resistant strains of HIV [17–21].

The World Health Organisation (WHO) commonly classifies HIVDR into three major categories; Acquired HIV Drug Resistance (ADR), Transmitted HIV Drug Resistance (TDR), and Pretreatment HIV Drug Resistance (PDR) [9]. Transmitted HIV drug resistance occurs when previously uninfected individuals are infected with a drug-resistant virus [22]. Acquired HIV drug resistance is Drug-Resistant Mutations (DRMs) that are selected during antiretroviral therapy, which can happen when viral replication is not fully suppressed in the drug's presence [23]. Pretreatment drug resistance refers to the resistance detected among ARV drug naïve people initiating ART or people with previous ARV drug exposure initiating or re-initiating first-line ART [9]. PDR is either TDR or ADR or both and it is associated with poor response to first-line treatment, premature virological failure, and accumulation of DRMs [9,24,25].

The development of anti-HIV vaccines is crucial in preventing HIV transmission and containing the HIV pandemic. However, the variability in the HIV-1 envelope proteins of more than 30% of the amino acids makes it difficult for anti-HIV vaccine development [8]. Currently, the development of broadly neutralizing anti-HIV antibodies (bNAbs) may help stimulate the immune response against HIV [26]. Vaccine candidates such as polyvalent envelope glycoprotein vaccines have shown greater success in the development of anti-HIV vaccines [27,28]. Despite much good progress made in anti-HIV vaccine development, no known vaccine has prevented HIV transmission to date [29].

The rise in HIV drug resistance is a great threat to global health, and if it is not urgently addressed, it may result in millions of deaths and an increase in new HIV infections. Therefore, the purpose of this narrative review was to review the published studies on HIV drug resistance to ART.

MATERIALS AND METHODS

A literature search for this narrative review was conducted from November 2019 to September 2020 with the following databases: PubMed, Google Scholar, and The Lancet. Keywords used in the search were 'antiretroviral therapy', 'resistance', 'HIV vaccines', 'HIV drug resistance', 'HIVDR', and the Boolean word 'AND'. Articles were read and assessed for relevance. The inclusion criteria included articles published from 1999 to 2021 and focused on HIV drug resistance. Exclusion criteria included editorials, commentaries, and conference abstracts. A total of 133 articles were retrieved but only 77 were included in this review. Five major themes were identified; mechanisms of HIV drug resistance, factors leading to HIV drug resistance, the prevalence of HIV drug resistance, the impact of HIV drug resistance on HIV/AIDS and strategies to mitigate HIV drug resistance.

Mechanisms of HIV drug resistance

Resistance occurs because of alterations called mutations in the HIV's genetic structure which lead to changes in certain proteins, most commonly enzymes that regulate the production of infectious viruses [30]. The mechanisms of resistance to antiretroviral drugs differ from one drug class to another [31]. The available classes of HIV drugs include Nucleoside Reverse Transcriptase Inhibitors (NRTIs), Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs), Protease Inhibitors (PIs), Integrase Inhibitors, Fusion Inhibitors, and CCR5 Antagonists [32]. Resistance to NRTIs occurs through modification of the reverse transcription enzyme allowing it to discriminate between NRTIs and analogue substrates which results in reduced incorporation of the NRTIs into the growing chain [32]. Another way in which resistance to NRTIs occurs is through enhanced removal of the drug from its site of attachment at the end of the Deoxyribonucleic Acid (DNA) chain [7]. The table 1 shows the major NRTI resistance mutations.

These mutations allow Adenosine Triphosphate (ATP) to bind to the active site which liberates the drug and terminates its effects [7]. The most common mutations of NRTIs include M184V, K65R and Thymidine Analogue Mutations (TAMs). The TAMs include; M41L, D67N, K219Q, and T215F [33].

NNRTIs resistance occurs due to single mutations around the reverse transcriptase pocket which decreases the binding of the drug [34]. The most common NNRTI mutations are K103N, G190A, Y181C, and Y188L [33]. Protease Inhibitors (PIs) resistance occurs primarily as a result of amino acid mutations that arise within or proximal to the catalytic binding site to the drug [7]. Integrase inhibitors resistance occurs due to mutations in residues surrounding the integrase active site. Fusion inhibitors mechanism of resistance occurs because of mutations in the gp41 transmembrane protein that interfere with the association of

Table 1: Major NRTI Resistance Mutations.

| | Non- TAMs | | | | TAMs | | | | |
|-------------|-----------|----|----|-----|------|----|----|-----|-----|
| | 184 | 65 | 74 | 115 | 41 | 67 | 70 | 215 | 219 |
| Cons | M | K | L | Y | M | D | K | T | K |
| 3TC | VI | R | | | | | | | |
| FTC | VI | R | | | | | | | |
| ABC | VI | R | VI | F | | | | | |
| TDF | | R | | | | | | | |
| ZDV | | | | | L | N | R | YF | Q |

Heptad Repeat 1 (HR1) and HR2 required for virus-cell fusion [35]. Recent resistance of HIV to Dolutegravir (DTG) HAS has been reported [36]. Table 2 shows the major resistance mutations to NNRTIs, PIs and integrase inhibitors.

Factors leading to HIV drug resistance

Understanding the interplay of factors that are associated with HIV drug resistance is an important factor, especially in Low to Middle-Income Countries (LMIC) [37]. In China, Xing and colleagues studied the incidence and factors associated with HIV drug resistance in patients. The independent factors that were associated with the development of resistance included initial ART regimen, ART distribution institution, a high baseline viral load (≥ 1000 copies/ml), and low baseline CD4 counts (0-199 cells/ μ l) [38]. A cross-sectional study by Zhou, et al. [39] showed that gender and treatment time were factors associated with the development of drug resistance. A comparison of the risk of drug resistance between patients treated for 1-2 years (RR = 13.616, 95% CI: 1.715-108.109), 2-3 years (RR = 19.556, 95% CI: 2.278-167.857), and more than 3 years (RR = 50.579, 95% CI: 4.855-526.891) was made and it was found that there was a greater possibility of HIV drug resistance emerging with longer treatment duration.

A study conducted by Ekong and colleagues on epidemiologic and viral predictors of antiretroviral drug resistance in Nigeria evaluated 159 cases with ADR and 299 cases without ADR. The factors that were associated with ADR included older age (age group 31-40) (OR = 2.35 [95%

CI: 1.29, 4.27], age group 41 + OR = 2.31 [95% CI: 1.11, 4.84], being unmarried (single) (OR = 0.40 [95% CI: 0.24-0.67], higher education level (secondary) (OR = 2.14 [95% CI: 1.11-4.13]), non adherence to care (OR = 2.48 [95% CI: 1.50-4.00]), long treatment duration (> 3 years) (OR = 1.80 [95% CI: 1.37-2.35]), lower CD4 count and a higher viral load (OR = 1.97 [95% CI: 1.44-2,54]) [40].

Ekong, et al. and Xing, et al. [38,40] both observed that low CD4 cell count and high viral load were associated with drug resistance development. Their results showed that patients with a low CD4 count were more likely to develop resistance. These results are consistent with Jose, et al. [41] whose results also demonstrated that resistance was more likely to occur in patients with a low CD4 count than those with a high cell count (> 350 cells/ μ L). High viral load was associated with resistance development because there is incomplete viral suppression in patients, which increases the risk of drug resistance [42]. Longer treatment duration (> 3 years) was found to be a risk factor of HIV drug resistance development by both Ekong *et al.* and Zhou *et al.* thus, showing that patients who have been on ART for a longer time are at risk of HIV developing resistance [39,40].

Ekong, et al. [40]. were able to associate more factors with the development of resistance because they conducted a case-control study between 2004-2011. Therefore, with the longer duration of the study, they were able to associate other factors with the development of drug resistance such as age, being unmarried and high education, non-

Table 2: Major NNRTIs, PIs and Integrase Inhibitor Resistance Mutations.

| | 47 | 50 | 54 | 82 | 92 | 101 | 103 | 106 | 132 | 181 | 188 | 190 |
|-------|----|----|------|------|----|-----|-----|-----|-----|-----|-----|-----|
| Cons | I | I | I | V | E | K | K | V | | Y | Y | G |
| NVP | | | | | | P | N | AM | | C | L | ASE |
| EFV | | | | | | P | N | AM | | | L | ASE |
| ETR | | | | | | P | | | | C | | EQ |
| RPV | | | | | | P | | | | C | L | EQ |
| ATV/r | | L | VTAM | ATSF | | | | | | | | |
| DRV/r | VA | V | LM | F | | | | | | | | |
| LPV/r | VA | V | VTAM | ATSF | | | | | | | | |
| RAL | | | | | Q | | | | KA | | | |
| EVG | | | | | Q | | | | KA | | | |

adherence. Being unmarried was associated with the risk of HIVDR development because the support that married individuals have makes them adhere to treatment more [40]. Higher education was associated with increased risk of drug resistance, but this differs from Boateng and Awunyo-Vitor who found that a high education level led to good knowledge on HIV/AIDS and ART, it influenced motivation and uptake of ART, thus making the patients more adherent and reducing the risk of HIV drug resistance [43]. Non-adherence to ART increases the possibility of HIV drug resistance and factors that are associated with poor adherence include gender, age, employment, marital status, and health status [44].

2.3. Stigma and social backgrounds among infected individuals are other factors that can lead to HIV drug resistance. This is because stigma and social backgrounds can affect access to ART and compliance [45]. A study conducted by Denison and colleagues found that self-stigma was associated with incomplete adherence among patients on ART. Due to self-stigma, some patients avoided attending nearby clinics for fear of being identified as HIV infected by the community and health care workers whom they know [46,47].

Prevalence of HIV drug resistance

The prevalence of HIVDR has been increasing over the past few years. Moreover, HIVDR has been reported among the first- and second-line ART [48]. This called for the third-line of treatment more especially in low- and middle-income countries [49]. In 1999, the prevalence of HIVDR was reported to be 16.3% among newly infected individuals sampled between 1994 and 1999 in the United States of America [12]. In 2010, the WHO reported the prevalence estimates for HIV drug resistance to NNRTIs to be 5.5% in Low and Middle-Income Countries (LMICs), and by 2015 Transmitted Drug Resistance (TDR) levels had risen in the ranges of 1-12.3% in different regions [22,50].

Transmitted Drug Resistance (TDR) levels have been rising over the past few years. A meta-analysis by Rhee, et al. [50] found the median overall TDR prevalence in sub-Saharan Africa and South/Southeast Asia to be 2.8% and 2.9% respectively and it was accompanied by an estimated 1.1-fold yearly increase in NRTI and NNRTI-associated TDR since the antiretroviral scale-up began. The TDR prevalence in upper-income Asian countries, Latin America/Caribbean, Europe, and North America was 5.6%, 7.6%, 9.4%, and 11.5% respectively with an estimated 1.1-fold yearly increase in TDR in North America and Latin America/Caribbean. In upper Asian countries, the estimated yearly increase of NNRTI associated TDR was 1.2-fold, while in Europe there was a 0.9-fold yearly decrease in NRTI-associated TDR. A study in Brazil also reported an increase in the overall TDR prevalence to be 16.3% (95% CI: 8.1-30.0) and the highest prevalence was against NNRTI (11.6%), with K103N being the most frequently identified mutation [51]. A 7.21% TDR

was reported recently in China in which HIV was resistant to NNRTI, NRTI, PIs, and PIs + NNRTI [52].

The findings from these studies on TDR show an increase in TDR with Brazil reporting the highest prevalence of 16.3%. The high prevalence rate reported in Brazil was because the study focused on patients with acute/recent HIV infection while the meta-analysis focused on recently and chronically infected ARV-naïve individuals. TDR studies focusing on chronically HIV-infected patients, usually have lower TDR detection because of the overgrowth of more fit, drug-sensitive viral quasi-species in such individuals [51].

The rise in TDR can at times be exceeded by a rise in acquired drug resistance as seen from a meta-analysis in China. The meta-analysis reported a pooled TDR prevalence of 3.0% including 0.3% for NRTIs, 1.4% for NNRTIs, and 0.5% for PIs, and a pooled ADR prevalence of 44.7% including 31.4% for NRTIs, 39.5% for NNRTIs, and 1.0% for PIs. The difference in HIVDR prevalence observed was regional due to the variable coverage rates and durations of ART since its introduction in China [53].

The prevalence of Pretreatment Drug Resistance (PDR) has been rising in many LMICs especially in sub-Saharan [54]. This rise in prevalence is high for the most commonly used first-line NNRTIs, namely efavirenz, and nevirapine. A meta-analysis done in 64 LMIC reported the estimated annual increase of PDR for NNRTIs of 23% in Southern Africa, 17% in Eastern Africa, 17% in Western and Central Africa, 11% in Latin America and the Caribbean, and 11% in Asia. The most common mutations reported in the study were K103N, Y181C, and G190A [55]. Similar findings for increased levels of prevalence of PDR for NNRTIs were reported in other studies. In a study done in West Africa and Southeast Asia, the recorded overall PDR prevalence was 15.9% (95% CI: 13.8-18.3) for any drug-resistant mutation, ranging from 9.6% and 10.2% in Burkina Faso and Thailand, respectively, 14.7% in Vietnam, 15.4% in Mali, 16.5% in Côte d'Ivoire and 19.3% in Cameroon, to 24.6% in Togo. The prevalence by ARV drug class was 3% for PI, 4% for NRTI, and 12% for NNRTI [56]. Bisso, et al. [57] from Argentina reported 14% PDR prevalence with 2%, 3%, and 11% prevalence for PIs, NRTIs, and NNRTIs respectively. The highest drug-specific prevalence was against NNRTIs then NRTIs and the most frequently associated mutations were M184V for NRTIs and K103N for NNRTIs. The high prevalence of NNRTIs observed is because in patients who have a virological failure the earliest mutations to emerge are NNRTI mutations and so transmission of these resistant variants is greater. The NRTI mutation does emerge early as well, but it is not commonly transmitted because of its significant fitness cost to the virus [55].

HIV drug resistance rates in African countries have also been reported. A study in sub-Saharan Africa on men who have sex with men (MSM) and transgender women found that 63 (34.4%) of 183 participants were infected with HIV.

Of the 63 participants, 11 (17.5%) had at least one major drug-resistant mutation, and the most frequently detected mutations were M184V for NRTI and K103N for NNRTI [58]. From Zimbabwe, Kouamou, et al. [59] detected HIV drug-resistant mutations in 97% (72/74) of the participants, with NNRTI drug-resistant mutations at 97%, and NRTI drug-resistant mutations at 84%. The most frequently identified mutations were M184V, K65R, and TAMs, also, two-class resistance was detected in 84% (62/74) of the participants owed to NRTIs and NNRTIs. In Kenya, the prevalence of drug resistance was reported to be 23.1% among the studied population with M184V and K103N been the most prevalent drug mutations [45]. Overall, Zimbabwe had the highest prevalence of HIV drug resistance. This could be because, in many LMICs, there is a lack of differentiation between patients who are being initiated on first-line ART and those who are being re-initiated on first-line ART [55]. The risk of HIV drug resistance is high in those who have previous exposure to ART [55].

Impact of HIV drug resistance on HIV/AIDS

Phillips, et al. [60] studied the impact of HIV drug resistance on HIV/AIDS and found that a high level of HIVDR indicates gaps in ART service delivery. Furthermore, the study brought to light that in situations where pretreatment drug resistance is over 10% the impact is great with an estimated 8,90,000 AIDS deaths and 45,000 new infections. This shows that HIV drug resistance causes attenuation of the potential full health benefits of ART and adds costs to the programs. Besides, the consequences of HIV drug resistance include treatment failure, the spread of drug-resistant HIV strains and increased mortality [60,61].

HIV drug resistance can cause ART failure, which may make the treatment of HIV more challenging [61,62]. It also leads to limitations in the choice of effective treatment regimens and increases costs switches to second and third-line therapy and routine laboratory monitoring of patients [20,63,64]. Furthermore, it can prolong the time required to achieve viral suppression and shorten the time to virological failure when compared to infections with a non- drug-resistant viral strain [65,66].

Strategies of mitigating HIV drug resistance

With the rise in NNRTI Pretreatment Drug Resistance (NNRTI PDR), WHO launched a Global Action Plan where it recommended that countries with high levels of NNRTI PDR defined by a PDR above 10% in populations starting ART, use a first-line regimen that remains active against HIV [67,68]. As of July 2018, WHO ART guidelines recommended the use of Dolutegravir (DTG)-based regimen as the preferred first-line antiretroviral therapy [68]. DTG is an integrase inhibitor with superior efficacy, tolerability and has a high genetic barrier to resistance than NNRTI, so the use of a DTG- based first-line regimen reduces the risk of HIV drug resistance [69]. The most commonly used DTG- based regimen is

Tenofovir Disoproxil Fumarate (TDF) + Lamivudine (3TC) + Dolutegravir (DTG) as first-line therapy [68,70,71].

Other ways of mitigating HIV drug resistance in patients on ART include increasing and improving access to viral load monitoring, enhancing HIV drug resistance testing services for individuals, improving adherence by providing person-centred care, enhancing research into long-acting ART formulations and functional care, encouraging ART programmes for implementation and promotion of good practices, increasing knowledge for better understanding of biological determinants of drug resistance, using ARVs with a high genetic barrier to resistance in first-line therapy [56,69,72,73].

The development of anti-HIV vaccines is urgently needed [74-77]. The HIV vaccines will prevent the new transmission and spread of HIV and HIV drug resistance. Despite a lot of challenges being faced in developing HIV vaccines, a breakthrough vaccine will change the disease burden across the globe.

CONCLUSION

HIV infections remain a global health problem that has been worsened by the development of HIV drug resistance. The current prevalence of HIV drug resistance is widespread in the world and it is a global threat to the treatment of HIV/AIDS. HIV drug resistance can cause attenuation of the potential full health benefits that people on ART can experience. Therefore, more research must be carried out to come up with various strategies like the use of ARVs with a high genetic barrier to resistance. This may help prevent the further spread of HIV drug resistance. Besides, HIVDR must be closely monitored so that treatment can be according to HIV susceptibility. Moreover, the development of HIV vaccines is urgently needed across the globe to contain the HIV pandemic.

ACKNOWLEDGEMENTS

We are grateful to the University of Zambia e-library for providing access to the articles used in this publication.

References

1. Iwuji C, Newell ML. Towards control of the global HIV epidemic: Addressing the middle-90 challenge in the UNAIDS 90-90-90 target. *PLoS Med.* 2017 May 2;14(5):e1002293. doi: 10.1371/journal.pmed.1002293. PMID: 28464003; PMCID: PMC5412986.
2. Piot P, Quinn TC. Response to the AIDS pandemic—a global health model. *N Engl J Med.* 2013 Jun 6;368(23):2210-8. doi: 10.1056/NEJMra1201533. Erratum in: *N Engl J Med.* 2013 Sep 19;369(12):1180. PMID: 23738546; PMCID: PMC3777557.
3. Nakazwe C, Michelo C, Sandøy IF, Fylkesnes K. Contrasting HIV prevalence trends among young women and men in Zambia in the past 12 years: data from demographic and health surveys 2002-2014. *BMC Infect Dis.* 2019 May 17;19(1):432. doi: 10.1186/s12879-019-4059-3. PMID: 31101081; PMCID: PMC6525340.
4. Aghokeng AF, Kouanfack C, Eymard-Duvernay S, Butel C, Edouard GE, Laurent C, Koulla-Shiro S, Delaporte E, Mpoudi-Ngole E, Peeters M. Virological outcome and patterns of HIV-1 drug resistance in patients with 36 months' antiretroviral therapy experience in Cameroon. *J Int AIDS Soc.* 2013 Jan 31;16(1):18004. doi: 10.7448/IAS.16.1.18004. PMID: 23374858; PMCID: PMC3562358.

5. Panos G, Samonis G, Alexiou VG, Kavarnou GA, Charatsis G, Falagas ME. Mortality and morbidity of HIV infected patients receiving HAART: a cohort study. *Curr HIV Res.* 2008 May;6(3):257-60. doi: 10.2174/157016208784324976. PMID: 18473789.
6. Krentz HB, Kliewer G, Gill MJ. Changing mortality rates and causes of death for HIV-infected individuals living in Southern Alberta, Canada from 1984 to 2003. *HIV Med.* 2005 Mar;6(2):99-106. doi: 10.1111/j.1468-1293.2005.00271.x. PMID: 15807715.
7. Sharma B. Drug Resistance in HIV-1: Genetic and Molecular Bases, Mechanisms, and Strategies to Combat the Issue. *Biochem Anal Biochem.* 2014;04(01):1-2. doi: 10.4172/2161-1009.1000e153
8. Gaschen B, Taylor J, Yusim K, Foley B, Gao F, Lang D, Novitsky V, Haynes B, Hahn BH, Bhattacharya T, Korber B. Diversity considerations in HIV-1 vaccine selection. *Science.* 2002 Jun 28;296(5577):2354-60. doi: 10.1126/science.1070441. PMID: 12089434.
9. WHO. HIV Drug Resistance Report 2019. Geneva, Switzerland. 2019. <https://bit.ly/3m39dro>
10. Jordan MR, Winsett J, Tiro A, Bau V, Barbara RS, Rowley C, Bellosillo N, Wanke C, Coakley EP. HIV Drug Resistance Profiles and Clinical Outcomes in Patients with Viremia Maintained at Very Low Levels. *World J AIDS.* 2013 Jun;3(2):71-78. doi: 10.4236/wja.2013.32010. PMID: 25664219; PMCID: PMC4319662.
11. Prosperi MC, Zazzi M, Punzi G, Monno L, Colao G, Corsi P, Di Giambenedetto S, Meini G, Ghisetti V, Bonora S, Pecorari M, Gismondo MR, Bagnarelli P, Carli T, De Luca A; ARCA Collaborative Group. Low rate of virological failure and maintenance of susceptibility to HIV-1 protease inhibitors with first-line lopinavir/ritonavir-based antiretroviral treatment in clinical practice. *J Med Virol.* 2010 Dec;82(12):1996-2003. doi: 10.1002/jmv.21927. PMID: 20981785.
12. Boden D, Hurlay A, Zhang L, Cao Y, Guo Y, Jones E, Tsay J, Ip J, Farthing C, Limoli K, Parkin N, Markowitz M. HIV-1 drug resistance in newly infected individuals. *JAMA.* 1999 Sep 22-29;282(12):1135-41. doi: 10.1001/jama.282.12.1135. PMID: 10501116.
13. UNAIDS JUNP on H. 90-90-90 an ambitious treatment target to help end the AIDS epidemic. UNAIDS Geneva, Switzerland; 2014. <https://bit.ly/3oiyBvW>
14. WHO. Meeting report on assessment of World Health Organization HIV drug resistance early warning indicators: report of the Early Advisory Indicator Panel meeting. 2012. <https://bit.ly/2Y2hbcw>
15. Dünya Sağlık Örgütü (DSÖ/WHO). Global Report on Early Warning Indicators of HIV Drug Resistance. World Health Organization. 2016;64. <https://bit.ly/3ijGD46>
16. Hamers RL, Schuurman R, Sigaloff KC, Wallis CL, Kityo C, Siwale M, Mandaliya K, Iwe P, Botes ME, Wellington M, Osibogun A, Wit FW, van Vugt M, Stevens WS, de Wit TF; PharmAccess African Studies to Evaluate Resistance (PASER) Investigators. Effect of pretreatment HIV-1 drug resistance on immunological, virological, and drug-resistance outcomes of first-line antiretroviral treatment in sub-Saharan Africa: a multicentre cohort study. *Lancet Infect Dis.* 2012 Apr;12(4):307-17. doi: 10.1016/S1473-3099(11)70255-9. Epub 2011 Oct 27. PMID: 22036233.
17. Xing H, Ruan Y, Li J, Shang H, Zhong P, Wang X, Liao L, Li H, Zhang M, Xue Y, Wang Z, Su B, Liu W, Dong Y, Ma Y, Li H, Qin G, Chen L, Pan X, Chen X, Peng G, Fu J, Chen RY, Kang L, Shao Y; Chinese National HIVDR Surveillance and Monitoring Network. HIV drug resistance and its impact on antiretroviral therapy in Chinese HIV-infected patients. *PLoS One.* 2013;8(2):e54917. doi: 10.1371/journal.pone.0054917. Epub 2013 Feb 6. PMID: 23405098; PMCID: PMC3566114.
18. Azam M, Malik A, Rizvi M, Rai A. Trends of drug-resistance-associated mutations in the reverse transcriptase gene of HIV type 1 isolates from North India. *Arch Virol.* 2014 Apr;159(4):719-25. doi: 10.1007/s00705-013-1889-y. Epub 2013 Oct 19. PMID: 24142273.
19. Wainberg MA, Zaharatos GJ, Brenner BG. Development of antiretroviral drug resistance. *N Engl J Med.* 2011 Aug 18;365(7):637-46. doi: 10.1056/NEJMr1004180. PMID: 21848464.
20. Sigaloff KC, Hamers RL, Wallis CL, Kityo C, Siwale M, Iwe P, Botes ME, Mandaliya K, Wellington M, Osibogun A, Stevens WS, van Vugt M, de Wit TF; PharmAccess African Studies to Evaluate Resistance (PASER). Unnecessary antiretroviral treatment switches and accumulation of HIV resistance mutations; two arguments for viral load monitoring in Africa. *J Acquir Immune Defic Syndr.* 2011 Sep 1;58(1):23-31. doi: 10.1097/QAI.0b013e318227fc34. PMID: 21694603.
21. Yashik S, Maurice M. Predicting a single HIV drug resistance measure from three international interpretation gold standards. *Asian Pac J Trop Med.* 2012 Jul;5(7):566-72. doi: 10.1016/S1995-7645(12)60100-X. PMID: 22647821.
22. WHO. HIV drug resistance report 2012. 2012. <https://bit.ly/3kRGS85>
23. Günthard HF, Calvez V, Paredes R, Pillay D, Shafer RW, Wensing AM, Jacobsen DM, Richman DD. Human Immunodeficiency Virus Drug Resistance: 2018 Recommendations of the International Antiviral Society-USA Panel. *Clin Infect Dis.* 2019 Jan 7;68(2):177-187. doi: 10.1093/cid/ciy463. PMID: 30052811; PMCID: PMC6321850.
24. Hofstra LM, Sauvageot N, Albert J, Alexiev I, Garcia F, Struck D, Van de Vijver DAMC, Åsjö B, Beshkov D, Coughlan S, Descamps D, Griskevicius A, Hamouda O, Horban A, Van Kasteren M, Kolupajeva T, Kostrikis LG, Liitsola K, Linka M, Mor O, Nielsen C, Otelea D, Paraskevis D, Paredes R, Poljak M, Puchhammer-Stöckl E, Sönnnerborg A, Staneková D, Stanojevic M, Van Laethem K, Zazzi M, Zidovec Lepej S, Boucher CAB, Schmit JC, Wensing AMJ; SPREAD Program, Puchhammer-Stöckl E, Sarcelletti M, Schmied B, Geit M, Balluch G, Vandamme AM, Vercauteren J, Derdelinckx I, Sasse A, Bogaert M, Ceunen H, De Roo A, De Wit S, Echahidi F, Fransens K, Goffard JC, Goubau P, Goudeseune E, Yombi JC, Lacor P, Liesnard C, Moutschen M, Pierard D, Rens R, Schrooten Y, Vaira D, Vandekerckhove LPR, Van den Heuvel A, Van Der Gucht B, Van Ranst M, Van Wijngaerden E, Vandercam B, Vekemans M, Verhofstede C, Clumeck N, Van Laethem K, Beshkov D, Alexiev I, Lepej SZ, Begovac J, Kostrikis L, Demetriades I, Kousiappa I, Demetriou V, Hezka J, Linka M, Maly M, Machala L, Nielsen C, Jørgensen LB, Gerstoft J, Mathiesen L, Parakeus D, Larsens A, Kvinesdal B, Liitsola K, Ristola M, Suni J, Sutinen J, Descamps D, Assoumou L, Castor G, Grude M, Flandre P, Storto A, Hamouda O, Kücherer C, Berg T, Braun P, Poggensee G, Däumer M, Eberle J, Heiken H, Kaiser R, Knechten H, Korn K, Müller H, Neifer S, Schmidt B, Walter H, Gussenheimer-Bartmeyer B, Harter T, Paraskevis D, Hatzakis A, Zavitsanou A, Vassilakis A, Lazanas M, Chini M, Lioni A, Sakka V, Kourkounti S, Pappazios V, Antoniadou A, Papadopoulos A, Poulakou G, Katsarolis I, Protopapas K, Chryssos G, Drimis S, Gargalianos P, Xylomenos G, Lourida G, Psychogiou M, Daikos GL, Sipsas NV, Kontos A, Gamaletsou MN, Koratzanis G, Sambatakou H, Mariolis H, Skoutelis A, Papastamopoulos V, Georgiou O, Panagopoulos P, Maltezos E, Coughlan S, De Gascun C, Byrne C, Duffy M, Bergin C, Reidy D, Farrell G, Lambert J, O'Connor E, Rochford A, Low J, Coakely P, O'Dea S, Hall W, Mor O, Levi I, Chemtob D, Grossman Z, Zazzi M, de Luca A, Balotta C, Riva C, Mussini C, Caramma I, Capetti A, Colombo MC, Rossi C, Prati F, Tramuto F, Vitale F, Ciccozzi M, Angarano G, Rezza G, Kolupajeva T, Vasin O, Griskevicius A, Lipnickiene V, Schmit JC, Struck D, Sauvageot N, Hemmer R, Arendt V, Michaux C, Staub T, Sequin-Devaux C, Wensing AMJ, Boucher CAB, van de Vijver DAMC, van Kessel A, van Bentum PHM, Brinkman K, Connell BJ, van der Ende ME, Hoepelman IM, van Kasteren M, Kuipers M, Langebeek N, Richter C, Santegoets RMWJ, Schrijnders-Gudde L, Schuurman R, van de Ven BJM, Åsjö B, Kran AB, Ormaasen V, Aavitsland P, Horban A, Stanczak JJ, Stanczak GP, Firlag-Burkacka E, Wiercinska-Drapalo A, Jablonowska E, Maolepsza E, Leszczyszyn-Pynka M, Szata W, Camacho R, Palma C, Borges F, Paixão T, Duque V, Araújo F, Otelea D, Paraschiv S, Tudor AM, Cernat R, Chiriac C, Dumitrescu F, Prisecariu LJ, Stanojevic M, Jevtovic D, Salemovic D, Staneková D, Habekova M, Chabadová Z, Drobkova T, Bukovinova P, Shunnar A, Truska P, Poljak M, Lunar M, Babic D, Tomazic J, Vidmar L, Vovko T, Karner P, Garcia F, Paredes R, Monge S, Moreno S, Del Amo J, Asensi V, Sirvent JL, de Mendoza C, Delgado R, Gutiérrez F, Berenguer J, Garcia-Bujalance S, Stella N, de Los Santos I, Blanco JR, Dalmou D, Rivero M, Segura F, Elías MJP, Alvarez M, Chueca N, Rodríguez-Martín C, Vidal C, Palomares JC, Viciana I, Viciana P, Cordoba J, Aguilera A, Domingo P, Galindo MJ, Miralles C, Del Pozo MA, Ribera E, Iribarren JA, Ruiz L, de la Torre J, Vidal F, Clotet B, Albert J, Heidarian A, Aperia-Peipke K, Axelsson M, Mild M, Karlsson A, Sönnnerborg A, Thalmé A, Navér L, Bratt G, Karlsson A, Blaxhult A, Gisslén M, Svennerholm B, Bergbrant I, Björkman P, Säll C, Mellgren Å, Lindholm A, Kuylenstierna N, Montelius R, Azimi F, Johansson B, Carlsson M, Johansson E, Ljungberg B, Ekvall H, Strand A, Mäkitalo S, Öberg S, Holmblad P, Höfer M, Holmberg H, Josefson P, Ryding U. Transmission of HIV Drug Resistance and the Predicted Effect on Current First-line Regimens in Europe. *Clin Infect Dis.* 2016 Mar 1;62(5):655-663. doi: 10.1093/cid/civ963. Epub 2015 Nov 29. PMID: 26620652; PMCID: PMC4741360.
25. Kityo C, Boerma RS, Sigaloff KCE, Kaudha E, Calis JCJ, Musiime V, Balinda S, Nakanjako R, Boender TS, Mugenyi PN, Rinke de Wit TF. Pretreatment HIV drug resistance results in virological failure and accumulation of additional resistance mutations in Ugandan children. *J Antimicrob Chemother.* 2017 Sep 1;72(9):2587-2595. doi: 10.1093/jac/dkx188. PMID: 28673027; PMCID: PMC5890670.
26. Ahmed Y, Tian M, Gao Y. Development of an anti-HIV vaccine eliciting broadly neutralizing antibodies. *AIDS Res Ther.* 2017 Sep 12;14(1):50. doi: 10.1186/s12981-017-0178-3. PMID: 28893278; PMCID: PMC5594608.
27. Cho MW, Kim YB, Lee MK, Gupta KC, Ross W, Plishka R, Buckler-White A, Igarashi T, Theodore T, Byrum R, Kemp C, Montefiori DC, Martin MA. Polyvalent envelope glycoprotein vaccine elicits a broader neutralizing antibody response but is unable to provide sterilizing protection against heterologous Simian/human immunodeficiency virus infection in pigtailed macaques. *J Virol.* 2001 Mar;75(5):2224-34. doi: 10.1128/JVI.75.5.2224-2234.2001. PMID: 11160726; PMCID: PMC114806.
28. Wang S, Pal R, Mascola JR, Chou TH, Mboudjeka I, Shen S, Liu Q, Whitney S, Keen T, Nair BC, Kalyanaraman VS, Markham P, Lu S. Polyvalent HIV-1 Env vaccine formulations delivered by the DNA priming plus protein boosting approach are effective in generating neutralizing antibodies against primary human immunodeficiency virus type 1 isolates from subtypes A, B, C, D and E. *Virology.* 2006 Jun 20;350(1):34-47. doi: 10.1016/j.virol.2006.02.032. Epub 2006 Apr 17. PMID: 16616287.
29. Haynes BF, Moody MA, Alam M, Bonsignori M, Verkoczy L, Ferrari G, Gao F, Tomaras GD, Liao HX, Kelsoe G. Progress in HIV-1 vaccine development. *J Allergy Clin*

- Immunol. 2014 Jul;134(1):3-10; quiz 11. doi: 10.1016/j.jaci.2014.04.025. Epub 2014 Jun 26. PMID: 25117798; PMCID: PMC4133697.
30. Tang MW, Liu TF, Shafer RW. The HIVdb system for HIV-1 genotypic resistance interpretation. *Intervirology*. 2012;55(2):98-101. doi: 10.1159/000331998. Epub 2012 Jan 24. PMID: 22286876; PMCID: PMC7068798.
 31. Asachop EL, Wainberg MA, Sloan RD, Tremblay CL. Antiviral drug resistance and the need for development of new HIV-1 reverse transcriptase inhibitors. *Antimicrob Agents Chemother*. 2012 Oct;56(10):5000-8. doi: 10.1128/AAC.00591-12. Epub 2012 Jun 25. PMID: 22733071; PMCID: PMC3457356.
 32. Haggblom A. Antiretroviral drug resistance and treatment outcomes of Human Immunodeficiency Virus type-1, implications for low and middle-income countries. Karolinska Institutet; 2016. <https://bit.ly/2Zx4hDu>
 33. Tang MW, Shafer RW. HIV-1 antiretroviral resistance: scientific principles and clinical applications. *Drugs*. 2012 Jun 18;72(9):e1-25. doi: 10.2165/11633630-000000000-00000. PMID: 22686620; PMCID: PMC3689909.
 34. Boucher CA, Bobkova MR, Geretti AM, Hung CC, Kaiser R, Marcelin AG, Streinu-Cercel A, van Wyk J, Dorr P, Vandamme AM. State of the Art in HIV Drug Resistance: Science and Technology Knowledge Gap. *AIDS Rev*. 2018 Jan-Mar;20(1):27-42. PMID: 29628515.
 35. Margeridon-Thermet S, Shafer RW. Comparison of the Mechanisms of Drug Resistance among HIV, Hepatitis B, and Hepatitis C. *Viruses*. 2010 Dec 1;2(12):2696-739. doi: 10.3390/v2122696. PMID: 21243082; PMCID: PMC3020796.
 36. Fokam J, Takou D, Semengue ENJ, Teto G, Beloumou G, Dambaya B, Santoro MM, Mossiang L, Billong SC, Cham F, Sosso SM, Temgoua ES, Nanfack AJ, Moudourou S, Kamgaing N, Kamgaing R, Ngako Pamen JN, Etame MMN, Bissek AZ, Elat JN, Moussi EE, Colizzi V, Perno CF, Ndjolo A; VIROFORUM. First case of Dolutegravir and Darunavir/r multi drug-resistant HIV-1 in Cameroon following exposure to Raltegravir: lessons and implications in the era of transition to Dolutegravir-based regimens. *Antimicrob Resist Infect Control*. 2020 Aug 26;9(1):143. doi: 10.1186/s13756-020-00799-2. PMID: 32843050; PMCID: PMC7449072.
 37. Rossouw TM, Feucht UD, Melikian G, van Dyk G, Thomas W, du Plessis NM, Avenant T. Factors Associated with the Development of Drug Resistance Mutations in HIV-1 Infected Children Failing Protease Inhibitor-Based Antiretroviral Therapy in South Africa. *PLoS One*. 2015 Jul 21;10(7):e0133452. doi: 10.1371/journal.pone.0133452. PMID: 26196688; PMCID: PMC4510388.
 38. Xing H, Wang X, Liao L, Ma Y, Su B, Fu J, He J, Chen L, Pan X, Dong Y, Liu W, Hsi JH, Yang L, Ruan Y, Shao Y. Incidence and associated factors of HIV drug resistance in Chinese HIV-infected patients receiving antiretroviral treatment. *PLoS One*. 2013 Apr 30;8(4):e62408. doi: 10.1371/journal.pone.0062408. PMID: 23638072; PMCID: PMC3640055.
 39. Zhou Y, Lu J, Wang J, Yan H, Li J, Xu X, Zhang Z, Qiu T, Ding P, Fu G, Huan X, Hu H. Prevalence of HIV Antiretroviral Drug Resistance and Its Impacts on HIV-1 Virological Failures in Jiangsu, China: A Cross-Sectional Study. *Biomed Res Int*. 2016;2016:1752437. doi: 10.1155/2016/1752437. Epub 2016 Oct 11. PMID: 27807537; PMCID: PMC5078637.
 40. Ekong E, Ndembi N, Okonkwo P, Dakum P, Idoko J, Banigbe B, Okuma J, Agaba P, Blattner W, Adebamowo C, Charurat M. Epidemiologic and viral predictors of antiretroviral drug resistance among persons living with HIV in a large treatment program in Nigeria. *AIDS Res Ther*. 2020 Feb 17;17(1):7. doi: 10.1186/s12981-020-0261-z. PMID: 32066473; PMCID: PMC7027291.
 41. Jose S, Quinn K, Dunn D, Cox A, Sabin C, Fidler S; UK CHIC and UK HDRD Steering Committees. Virological failure and development of new resistance mutations according to CD4 count at combination antiretroviral therapy initiation. *HIV Med*. 2016 May;17(5):368-72. doi: 10.1111/hiv.12302. Epub 2015 Aug 25. PMID: 26306942; PMCID: PMC4949527.
 42. Crowell TA, Pinyakorn S, Sacdalan C, Kroon E, Colby DJ, Puttamaswin S, Ubolyam S, Trichavaroj R, Butterworth O, Turk E, McCullough C, Chomont N, de Souza M, Robb ML, Phanuphak N, Ananworanich J; RV254/SEARCH010 Study Group. Viral Blips After Treatment Initiation During Acute Human Immunodeficiency Virus Infection. *Clin Infect Dis*. 2020 Jun 10;70(12):2706-2709. doi: 10.1093/cid/ciz936. PMID: 31550044; PMCID: PMC7286371.
 43. Boateng D, Awunyo-Vitor D. Knowledge, Perceptions, and Practices on Antiretroviral Therapy in Farming Communities in Ghana: A Study of HIV Positive Women. *Public Heal Res*. 2012;2(5):136-142.
 44. Ibrahim Y, Sutan R, Bin K, Latif A, Amara A, Adam I. Poor Adherence to Antiretroviral Therapy and associated factors among people living with HIV in Omdurman city, Sudan. *Malaysian J Public Heal Med*. 2014;14(1):90-101. <https://bit.ly/3CZ1fpv>
 45. Kinyua JG, Lihana RW, Kiptoo M, Muasya T, Odera I, Muiruri P, Songok EM. Antiretroviral resistance among HIV-1 patients on first-line therapy attending a comprehensive care clinic in Kenyatta National Hospital, Kenya: a retrospective analysis. *Pan Afr Med J*. 2018 Apr 2;29:186. doi: 10.11604/pamj.2018.29.186.10796. PMID: 30061964; PMCID: PMC6061825.
 46. Denison JA, Koole O, Tsui S, Menten J, Torpey K, van Praag E, Mukadi YD, Colebunders R, Auld AF, Agolory S, Kaplan JE, Mulenga M, Kwesigabo GP, Wabwire-Mangen F, Bangsberg DR. Incomplete adherence among treatment-experienced adults on antiretroviral therapy in Tanzania, Uganda and Zambia. *AIDS*. 2015 Jan 28;29(3):361-71. doi: 10.1097/QAD.0000000000000543. PMID: 25686684; PMCID: PMC4728173.
 47. Qiao S, Zhang Y, Li X, Menon JA. Facilitators and barriers for HIV-testing in Zambia: A systematic review of multi-level factors. *PLoS ONE*. 2018;13:e0192327. <https://bit.ly/3zTi5Vg>
 48. Hosseinipour MC, Gupta RK, Van Zyl G, Eron JJ, Nachega JB. Emergence of HIV drug resistance during first- and second-line antiretroviral therapy in resource-limited settings. *J Infect Dis*. 2013 Jun 15;207 Suppl 2(Suppl 2):S49-56. doi: 10.1093/infdis/jit107. PMID: 23687289; PMCID: PMC3708738.
 49. Grinsztejn B, Hughes MD, Ritz J, Salata R, Mugenyi P, Hogg E, Wiclaw L, Gross R, Godfrey C, Cardoso SW, Bukuru A, Makanga M, Faesen S, Mave V, Wangari Ndege B, Nerette Fontain S, Samaneka W, Secours R, van Schalkwyk M, Mngqibisa R, Mohapi L, Valencia J, Sugandhavesa P, Montalban E, Avihingsanon A, Santos BR, Kumarasamy N, Kanyama C, Schooley RT, Mellors JW, Wallis CL, Collier AC; A5288 Team. Third-line antiretroviral therapy in low-income and middle-income countries (ACTG A5288): a prospective strategy study. *Lancet HIV*. 2019 Sep;6(9):e588-e600. doi: 10.1016/S2352-3018(19)30146-8. Epub 2019 Jul 29. PMID: 31371262; PMCID: PMC6857629.
 50. Rhee SY, Blanco JL, Jordan MR, Taylor J, Lemey P, Varghese V, Hamers RL, Bertagnolio S, Rinke de Wit TF, Aghokeng AF, Albert J, Avi R, Avila-Rios S, Bessong PO, Brooks JI, Boucher CA, Brumme ZL, Busch MP, Bussmann H, Chaix ML, Chin BS, D'Aquin TT, De Gascun CF, Derache A, Descamps D, Deshpande AK, Djoko CF, Eshleman SH, Fleury H, Frange P, Fujisaki S, Harrigan PR, Hattori J, Holguin A, Hunt GM, Ichimura H, Kaleebu P, Katzenstein D, Kiertburanakul S, Kim JH, Kim SS, Li Y, Lutsar I, Morris L, Ndembi N, Ng KP, Paranjape RS, Peeters M, Poljak M, Price MA, Ragonnet-Cronin ML, Reyes-Terán G, Rolland M, Sirivichayakul S, Smith DM, Soares MA, Soriano VV, Ssemwanga D, Stanojevic M, Stefani MA, Sugiura W, Sungkanupparh S, Tanuri A, Tee KK, Truong HM, van de Vijver DA, Vidal N, Yang C, Yang R, Yebra G, Ioannidis JP, Vandamme AM, Shafer RW. Geographic and temporal trends in the molecular epidemiology and genetic mechanisms of transmitted HIV-1 drug resistance: an individual-patient- and sequence-level meta-analysis. *PLoS Med*. 2015 Apr 7;12(4):e1001810. doi: 10.1371/journal.pmed.1001810. Erratum in: *PLoS Med*. 2015 Jun;12(6):e1001845. PMID: 25849352; PMCID: PMC4388826.
 51. Ferreira ACG, Coelho LE, Grinsztejn E, Jesus CS, Guimarães ML, Veloso VG, Grinsztejn B, Cardoso SW. Transmitted drug resistance in patients with acute/recent HIV infection in Brazil. *Braz J Infect Dis*. 2017 Jul-Aug;21(4):396-401. doi: 10.1016/j.bjid.2017.03.013. Epub 2017 May 21. PMID: 28539254.
 52. Pang X, Tang K, He Q, Huang J, Fang N, Zhou X, Zhu Q, Wu X, Shen Z, Liang S. HIV drug resistance and HIV transmission risk factors among newly diagnosed individuals in Southwest China. *BMC Infect Dis*. 2021 Feb 8;21(1):160. doi: 10.1186/s12879-021-05854-w. PMID: 33557775; PMCID: PMC7871613.
 53. Zuo L, Liu K, Liu H, Hu Y, Zhang Z, Qin J, Xu Q, Peng K, Jin X, Wang JH, Zhang C. Trend of HIV-1 drug resistance in China: A systematic review and meta-analysis of data accumulated over 17 years (2001-2017). *EClinicalMedicine*. 2020 Jan 5;18:100238. doi: 10.1016/j.eclinm.2019.100238. Erratum in: *EClinicalMedicine*. 2021 Mar 12;33:100696. PMID: 31922125; PMCID: PMC6948268.
 54. Gupta RK, Jordan MR, Sultan BJ, Hill A, Davis DH, Gregson J, Sawyer AW, Hamers RL, Ndembi N, Pillay D, Bertagnolio S. Global trends in antiretroviral resistance in treatment-naïve individuals with HIV after rollout of antiretroviral treatment in resource-limited settings: a global collaborative study and meta-regression analysis. *Lancet*. 2012 Oct 6;380(9849):1250-8. doi: 10.1016/S0140-6736(12)61038-1. Epub 2012 Jul 23. Erratum in: *Lancet*. 2012 Oct 6;380(9849):1230. PMID: 22828485; PMCID: PMC3790969.
 55. Gupta RK, Gregson J, Parkin N, Haile-Selassie H, Tanuri A, Andrade Forero L, Kaleebu P, Wataha C, Aghokeng A, Mutenda N, Dzangare J, Hone S, Hang ZZ, Garcia J, Garcia Z, Marchorro P, Beteta E, Giron A, Hamers R, Inzaule S, Frenkel LM, Chung MH, de Oliveira T, Pillay D, Naidoo K, Kharsany A, Kugathasan R, Cutino T, Hunt G, Avila Rios S, Doherty M, Jordan MR, Bertagnolio S. HIV-1 drug resistance before initiation or re-initiation of first-line antiretroviral therapy in low-income and middle-income countries: a systematic review and meta-regression analysis. *Lancet Infect Dis*. 2018 Mar;18(3):346-355. doi: 10.1016/S1473-3099(17)30702-8. Epub 2017 Dec 5. Erratum in: *Lancet Infect Dis*. 2017 Dec 15; PMID: 29198909; PMCID: PMC5835664.
 56. Ngo-Giang-Huong N, Huynh THK, Dagnra AY, Toni TD, Maiga AI, Kania D, Eymard-Duvernay S, Peeters M, Soulie C, Peytavin G, Rekacewicz C, Chaix ML, Aghokeng AF; ANRS 12333 Study Group. Prevalence of pretreatment HIV drug resistance in West African and Southeast Asian countries. *J Antimicrob Chemother*. 2019 Feb 1;74(2):462-467. doi: 10.1093/jac/dky443. PMID: 30418575; PMCID: PMC6337899.

57. Bissio E, Barbás MG, Bouzas MB, Cudolá A, Salomón H, Espínola L, Fernández Giuliano S, Kademian S, Mammama L, Ornani ML, Ravasi G, Vila M, Zapiola I, Falistocco C. Pretreatment HIV-1 drug resistance in Argentina: results from a surveillance study performed according to WHO-proposed new methodology in 2014-15. *J Antimicrob Chemother.* 2017 Feb;72(2):504-510. doi: 10.1093/jac/dkw445. Epub 2016 Oct 26. PMID: 27789684.
58. Zhang Y, Fogel JM, Guo X, Clarke W, Breud A, Cummings V, Hamilton EL, Ogendo A, Kayange N, Panchia R, Dominguez K, Chen YQ, Sandfort T, Eshleman SH. Antiretroviral drug use and HIV drug resistance among MSM and transgender women in sub-Saharan Africa. *AIDS.* 2018 Jun 19;32(10):1301-1306. doi: 10.1097/QAD.0000000000001839. PMID: 29794492; PMCID: PMC6146920.
59. Kouamou V, Varyani B, Shamu T, Mapangisana T, Chimbetete C, Mudzviti T, Manasa J, Katzenstein D. Drug Resistance Among Adolescents and Young Adults with Virologic Failure of First-Line Antiretroviral Therapy and Response to Second-Line Treatment. *AIDS Res Hum Retroviruses.* 2020 Jul;36(7):566-573. doi: 10.1089/AID.2019.0232. Epub 2020 Apr 2. PMID: 32138527.
60. Phillips AN, Stover J, Cambiano V, Nakagawa F, Jordan MR, Pillay D, Doherty M, Revill P, Bertagnolio S. Impact of HIV Drug Resistance on HIV/AIDS-Associated Mortality, New Infections, and Antiretroviral Therapy Program Costs in Sub-Saharan Africa. *J Infect Dis.* 2017 May 1;215(9):1362-1365. doi: 10.1093/infdis/jix089. Erratum in: *J Infect Dis.* 2017 Nov 15;216(8):1048. PMID: 28329236; PMCID: PMC5451603.
61. Cambiano V, Bertagnolio S, Jordan MR, Lundgren JD, Phillips A. Transmission of drug resistant HIV and its potential impact on mortality and treatment outcomes in resource-limited settings. *J Infect Dis.* 2013 Jun 15;207 Suppl 2:S57-62. doi: 10.1093/infdis/jit111. PMID: 23687290.
62. Luo XL, Mo LD, Su GS, Huang JP, Wu JY, Su HZ, Huang WH, Luo SD, Ni ZY. Incidence and types of HIV-1 drug resistance mutation among patients failing first-line antiretroviral therapy. *J Pharmacol Sci.* 2019 Apr;139(4):275-279. doi: 10.1016/j.jpshs.2018.11.016. Epub 2019 Jan 17. PMID: 30928089.
63. Boender TS, Hoenderboom BM, Sigaloff KC, Hamers RL, Wellington M, Shamu T, Siwale M, Labib Maksimos EE, Nankya I, Kityo CM, Adeyemo TA, Akanmu AS, Mandaliya K, Botes ME, Ondoa P, Rinke de Wit TF. Pretreatment HIV drug resistance increases regimen switches in sub-Saharan Africa. *Clin Infect Dis.* 2015 Dec 1;61(11):1749-58. doi: 10.1093/cid/civ656. Epub 2015 Aug 3. PMID: 26240203.
64. Petersen ML, Tran L, Geng EH, Reynolds SJ, Kambugu A, Wood R, Bangsberg DR, Yiannoutsos CT, Deeks SG, Martin JN. Delayed switch of antiretroviral therapy after virologic failure associated with elevated mortality among HIV-infected adults in Africa. *AIDS.* 2014 Sep 10;28(14):2097-107. doi: 10.1097/QAD.0000000000000349. PMID: 24977440; PMCID: PMC4317283.
65. De Beaudrap P, Thiam M, Diouf A, Toure-Kane C, Ngom-Guèye NF, Vidal N, Mboup S, Ndoye I, Sow PS, Delaporte E; ANRS 1215 Study Group. Risk of virological failure and drug resistance during first and second-line antiretroviral therapy in a 10-year cohort in Senegal: results from the ANRS 1215 cohort. *J Acquir Immune Defic Syndr.* 2013 Apr 1;62(4):381-7. doi: 10.1097/QAI.0b013e31827a2a7a. PMID: 23117504.
66. Hamers RL, Sigaloff KC, Kityo C, Mugenyi P, de Wit TF. Emerging HIV-1 drug resistance after roll-out of antiretroviral therapy in sub-Saharan Africa. *Curr Opin HIV AIDS.* 2013 Jan;8(1):19-26. doi: 10.1097/COH.0b013e32835b7f94. PMID: 23143140.
67. WHO. Guidelines on the public health response to pretreatment HIV drug resistance. 2017. <https://bit.ly/3D0IEez>
68. WHO. Updated recommendations on first-line and second-line antiretroviral regimens and post-exposure prophylaxis and recommendations on early infant diagnosis of HIV. 2018. <https://bit.ly/3m5HaaV>
69. Kanters S, Vitoria M, Doherty M, Socias ME, Ford N, Forrest JI, Popoff E, Bansback N, Nsanzimana S, Thorlund K, Mills EJ. Comparative efficacy and safety of first-line antiretroviral therapy for the treatment of HIV infection: a systematic review and network meta-analysis. *Lancet HIV.* 2016 Nov;3(11):e510-e520. doi: 10.1016/S2352-3018(16)30091-1. Epub 2016 Sep 6. PMID: 27658869.
70. Meintjes G, Moorhouse MA, Carmona S, Davies N, Dlamini S, van Vuuren C, Manzini T, Mathe M, Moosa Y, Nash J, Nel J, Pakade Y, Woods J, Van Zyl G, Conradie F, Venter F. Adult antiretroviral therapy guidelines 2017. *South Afr J HIV Med.* 2017 Jul 15;18(1):776. doi: 10.4102/sajhivmed.v18i1.776. PMID: 29568644; PMCID: PMC5843236.
71. Amstutz A, Nsakala BL, Vanobberghen F, Muhairwe J, Glass TR, Namane T, Mpholo T, Battegay M, Klimkait T, Labhardt ND. Switch to second-line versus continued first-line antiretroviral therapy for patients with low-level HIV-1 viremia: An open-label randomized controlled trial in Lesotho. *PLoS Med.* 2020 Sep 16;17(9):e1003325. doi: 10.1371/journal.pmed.1003325. PMID: 32936795; PMCID: PMC7494118.
72. Boyer S, March L, Kouanfack C, Laborde-Balen G, Marino P, Aghokeng AF, Mpoudi-Ngole E, Koulla-Shiro S, Delaporte E, Carrieri MP, Spire B, Laurent C, Moatti JP; Stratall ANRS 12110/ESTHER Study Group. Monitoring of HIV viral load, CD4 cell count, and clinical assessment versus clinical monitoring alone for antiretroviral therapy in low-resource settings (Stratall ANRS 12110/ESTHER): a cost-effectiveness analysis. *Lancet Infect Dis.* 2013 Jul;13(7):577-86. doi: 10.1016/S1473-3099(13)70073-2. Epub 2013 Apr 18. PMID: 23602084.
73. Gaolathe T, Wirth KE, Holme MP, Makhema J, Moyo S, Chakalisa U, Yankinda EK, Lei Q, Mmalane M, Novitsky V, Okui L, van Widenfelt E, Powis KM, Khan N, Bennett K, Bussmann H, Dryden-Peterson S, Lebelonyane R, El-Halabi S, Mills LA, Marukutira T, Wang R, Tchetgen EJ, DeGruttola V, Essex M, Lockman S; Botswana Combination Prevention Project study team. Botswana's progress toward achieving the 2020 UNAIDS 90-90-90 antiretroviral therapy and virological suppression goals: a population-based survey. *Lancet HIV.* 2016 May;3(5):e221-30. doi: 10.1016/S2352-3018(16)00037-0. Epub 2016 Mar 24. PMID: 27126489; PMCID: PMC5146754.
74. Ensofi B, Cafaro A, Monini P, Marcotullio S, Ensofi F. Challenges in HIV Vaccine Research for Treatment and Prevention. *Front Immunol.* 2014 Sep 8;5:417. doi: 10.3389/fimmu.2014.00417. PMID: 25250026; PMCID: PMC4157563.
75. Ng'uni T, Chasara C, Ndhlovu ZM. Major Scientific Hurdles in HIV Vaccine Development: Historical Perspective and Future Directions. *Front Immunol.* 2020 Oct 28;11:590780. doi: 10.3389/fimmu.2020.590780. PMID: 33193428; PMCID: PMC7655734.
76. Esparza J. A brief history of the global effort to develop a preventive HIV vaccine. *Vaccine.* 2013 Aug 2;31(35):3502-18. doi: 10.1016/j.vaccine.2013.05.018. Epub 2013 May 21. PMID: 23707164.
77. Phillips AN, Cambiano V, Nakagawa F, Ford D, Lundgren JD, Roset-Bahmanyar E, Roman F, Van Effelterre T. Potential future impact of a partially effective HIV vaccine in a southern African setting. *PLoS One.* 2014 Sep 10;9(9):e107214. doi: 10.1371/journal.pone.0107214. PMID: 25207973; PMCID: PMC4160197.

How to cite this article: Phiri MN, Mudenda S. Human Immunodeficiency Virus (HIV) Drug Resistance: A Global Narrative Review. *J Biomed Res Environ Sci.* 2021 Sept 30; 2(9): 857-864. doi: 10.37871/jbres1323, Article ID: JBRES1323, Available at: <https://www.jelsciences.com/articles/jbres1323.pdf>